Echinaforce®

SUGGESTED USE

Adults add 15-20 drops to a small amount of water 3 to 5 times daily before eating. Salivate well before swallowing.

Tablets: 1-2 tablets 3-5 times a day. Let the tablets dissolve in mouth.

Children: 5 drops in a small amount of water 3 times daily before eating.

When used as a preventative, dosage should be repeated for five consecutive days then halted for two days. This cycle should be repeated until desired results are observed. In acute cases, 40 drops 3 times daily may be used for up to 3 days. It is suggested the product not be used for more than 2 months continuously.

Tolerance

Echinaforce is tolerated very well. Hypersensitivity reactions (for example, skin rashes) have not appeared with Echinaforce to date. Should such reactions occur discontinue use immediately and consult a physician. Echinacea products should be avoided by persons with known allergy to plants of the Composite family, or individuals with immune system disorders.

Supports a healthy immune system and natural resistance during winter season

- 100% organically certified
- efficacy supported by clinical study
- unique combination of fresh Echinacea purpurea herb and root

Ingredients

Extract of fresh organically grown Echinacea herb, 95% herb and 5% roots. Sixty-seven percent (65%) alcohol is used to extract the maximum number and levels of active and supportive elements. 20 drops is equivalent to 259 mg fresh Echinacea. 2 tablets equivalent to 538 mg fresh Echinacea. The tablets contain approximately 250 mg lactose. Echinaforce has a full spectrum potency, meaning that the chemical profile of the extract is very similar to the whole fresh plant.

1 bottle 1.7 fl. oz. provides a 35 day supply - 1 bottle 3.4 fl. oz. provides a 70 day supply - 1 bottle 120 tabs provides a 40 day supply.

Constituents and Mode of Action

A broad range of investigations have shown that echinacea extracts stimulate



Packaging

Bottles 1.7 fl. oz./50 ml.
Bottles 3.4 fl. oz./100 ml.
Bottles of 120 tablets.



the nonspecific resistance system. The following primary effects of echinacea preparations have been observed in vivo and/or vitro:

By stimulating the bone marrow, they induce an increased release of leucocytes, especially granulocytes, into the blood. This promotes the resistance reaction.

The phagocytic potency of the granulocytes and macrophages is significantly increased. The serum properdin level is increased, promoting the non specific resistance system.

- The macrophages are stimulated to release effector substances such as tumor necrosis factor a, interferon b2, and interleukin 1.
- The antiphlogistic effect and the inhibition of cell processes leading to cross-linking the collagen also supports wound healing.

Even though many ingredients and elements of Echinacea have been isolated and identified, it is not possible to ascribe the effectiveness clearly to a fixed substance. Among the best known active constituents of Echinacea are:

- cichoric acid
- alkylamides
- coffeeaciderative
- polysaccharide
- glycoproteins
- flavonoids
- essential oil

Keeping in mind these ingredietns, we are convinced that an extract of the entire plant (all active and supporting ingredients) provides you with the most beneficial Echinacea product available.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Manufacturer * Bioforce AG * 9325 Roggwil TG * Switzerland
Exclusive Distributor * Bioforce USA * Chatham, NY 12037 * 800-641-7555





webmaster - Winnovative Technology Consulting, LLC