

- Aged Garlic Extract™ [300 mg]; Lecithin [190 mg] (per capsule).
- Aged Garlic Extract supports healthy cholesterol levels[1-17] and healthy nerve function,[18-23] and may quench free radicals that may oxidize LDL cholesterol and damage heart tissue.[24-26]
- Lecithin and its main constituent, choline, are important building blocks for all cell membranes.[27] Choline is necessary to make the crucial neurotransmitter, acetylcholine, in the brain and nerves.[28]
- Studies have suggested that choline and lecithin may help to maintain healthy cholesterol levels and nerves.[29-34]
- Garlic combined with lecithin has been shown to provide nutritional support for healthy cholesterol and blood pressure and to support healthy circulation.[35]
- **Supplement Facts**
- **Scientific References**
- **Next Kyolic Product**
- **Previous Kyolic Product**
- **All Kyolic Products**
- **Consumer Price List**

Kyolic Formula 104

SUGGESTED USE:
Take two capsules **with a meal** twice daily.

Supplement Facts Serving Size 2 capsules		**Product Codes:** **104-41: 100 capsules** **104-42: 200 capsules**
	Amount per serving	%Daily Value
Aged Garlic Extract™†† Powder (bulb)	600mg	**
Lecithin	380mg	**
††Special Garlic Preparation **Daily Value not established		

OTHER INGREDIENTS:
Gelatin, Silica and Magnesium Stearate (vegetable source).

Yeast, Sodium, Gluten, Sugar, Preservatives and Dairy Free.

TOP

REFERENCES FOR KYOLIC FORMULA 104

1. Abuirmeileh, N. et al. FASEB J. 5(6): 8048, 1991.
2. Efendy, J. et al. J. Vasc. Res. 33(S1): 23, 1996.
3. Efendy, J. et al. Atherosclerosis 132: 37-42, 1997.
4. Kawashima, Y. et al. Shinryo To Shinyaku (Treat. New Med.) 26: 377-388, 1989.
5. Lau, B. et al. Nutr. Res. 7: 139-149, 1987.
6. Liu, L. and Yeh, Y. 1999. FASEB 13(4): A556; #442.7.
7. Liu, L. and Yeh, Y.Y. 1998. FASEB A261:1523.
8. Qureshi, A. et al. First World Congress on the Health Significance of Garlic and Garlic Constituents (FWC). Washington, D.C. August 28-30, p. 16, 1990.
9. Steiner, M. and Lin, R. 1994. J. Amer. Coll. Nutr 13(5): 524.
10. Steiner, M. et al. Am. J. Clin. Nutr. 64: 866-870, 1996.
11. Steiner, M. et al. J. Am. Coll. Nutr. 13(5): 524, 1994.
12. Steiner, M . et al. Shinyaku To Rinsho (New Drug Clin.) 45(3): 456-466, 1996.
13. Yeh, Y. et al. J. Am. Coll. Nutr. 14(5):545, 1995.
14. Yeh, Y. et al. FWC. Washington, D.C. August 28-30, p. 37, 1990.
15. Yeh, Y. et al. Lipids 29(3): 189-193, 1994.
16. Yeh, Y. Shinyaku To Rinsho (New Drug Clin.) 45(3): 441-450, 1996.
17. Yeh, Y. et al. In: Food Factors for Cancer Prevention. Ohigashi, et al. eds. Springer-ag, Tokyo, p 226-230, 1997.
18. Welch, C. et al. Cancer Lett. 63: 211-219, 1992.
19. Gwebu, E. et al. Soc. Neurosci. 24(2):#571.11, 1998.
20. Moriguchi, T. et al. Neurochem. Res. 22(12): 1449-1452, 1997.
21. Moriguchi, T. et al. Phytother. Res. 10: 468-472, 1996.
22. Nishiyama, N. et al. Int. Acad. Biomed. Drug Res. Basel. Karger. 11: 253-258, 1996.
23. Saito, H. et al. Ch. 16. In: Nutraceuticals: Designer Foods III Garlic, Soy and Licorice. Paul P. Lanchance (ed.). Food & Nutrition Press, Trumbell, CT, pp. 153-161, 1997.
24. Steiner, M. et al. J. Cardiovascular Pharmacol. 31: 904-908, 1998.
25. Mundy, J. et al. Atherosclerosis 143: 399-404, 1999.
26. Ide, N. & Lau, B. J. Pharm. Pharmacol. 49: 908-911, 1997.
27. Recommended Dietary Allowances (RDA). 10th ed. National Academy Press, Washington, D.C., 1989, pp.263-264.
28. Cohen, E. & Wurtman, R. Science 191: 561-562, 1976.
29. Sitaram, N. et al. Science 201: 274-276, 1978.
30. Sitaram, N. et al. Life Science 22: 1555-1560, 1978.
31. Linscheer, W. and Vergroesen, A. In Modern Nutrition in Health and Disease. (Shils, M.E. and Young, V.R. Ed.) 7th Edition. Philadelphia, Lea & Febiger, 1988, p.91.

32. O'Mullane, J. et al. Atherosclerosis 45: 81-90, 1982.
33. Beil, F. et al. J. Lipid Res. 21: 525-536, 1980.
34. Boyd, W. et al. Lancet 2: 711, 1977.
35. Samochowiec, L. & Wojcicki, J. Arztliche Forschung 37: 1-8, 1990.

