

- Aged Garlic Extract™ [300 mg]; Whey (per capsule/tablet).
- Aged Garlic supports healthy cholesterol levels,[1-17] circulation,[18-21] immune function,[21-32] liver function,[41-48] and nerves[49-54] and assists in fighting stress and fatigue.[33-40]
- Aged Garlic Extract and its constituents have also been shown to enhance the growth of beneficial bacteria.[57-58]
- Whey contains a high quality protein[59] and is an excellent food source for the growth of beneficial bacteria.[60]
- **Supplement Facts**
- **Scientific References**
- **Next Kyolic Product**
- **All Kyolic Products**
- **Consumer Price List**

Kyolic Formula 100

SUGGESTED USE:
Take two tablets/capsules **with a meal** twice daily.

Supplement Facts Serving Size 2 tablets/capsules		**Product Codes:** **100-31: 100 tablets** **100-32: 200 tablets** **100-41: 100 capsules** **100-42: 200 capsules** **100-43: 300 capsules**
	Amount per serving	%Daily Value
Aged Garlic Extract†† Powder (bulb)	600mg	**
††Special Garlic Preparation **Daily Value not established.		

OTHER TABLET INGREDIENTS:
Whey, Alginic acid (seaweed), Silica, Cellulose, and Magnesium Stearate (vegetable source).

OTHER CAPSULE INGREDIENTS:
Gelatin, Whey and Magnesium Stearate (vegetable source).

Yeast, Sodium, Preservatives, Sugar, and Gluten Free.

TOP

REFERENCES FOR KYOLIC FORMULA 100

1. Abuirmeileh, N. et al. FASEB J. 5(6): 8048, 1991.
2. Efendy, J. et al. J. Vasc. Res. 33(S1): 23, 1996.
3. Efendy, J. et al. Atherosclerosis 132: 37-42, 1997.
4. Kawashima, Y. et al. Shinryo To Shinyaku (Treat. New Med.) 26: 377-388, 1989.
5. Lau, B. et al. Nutr. Res. 7: 139-149, 1987.
6. Liu, L. and Yeh, Y. 1999. FASEB 13(4): A556; #442.7.
7. Liu, L. and Yeh, Y.Y. 1998. FASEB A261:1523.
8. Qureshi, A. et al. First World Congress on the Health Significance of Garlic and Garlic Constituents (FWC). Washington, D.C. August 28-30, p. 16, 1990.
9. Steiner, M. and Lin, R. 1994. J. Amer. Coll. Nutr 13(5): 524.
10. Steiner, M. et al. Am. J. Clin. Nutr. 64: 866-870, 1996.
11. Steiner, M. et al. J. Am. Coll. Nutr. 13(5): 524, 1994.
12. Steiner, M . et al. Shinyaku To Rinsho (New Drug Clin.) 45(3): 456-466, 1996.
13. Yeh, Y. et al. J. Am. Coll. Nutr. 14(5):545, 1995.
14. Yeh, Y. et al. FWC. Washington, D.C. August 28-30, p. 37, 1990.
15. Yeh, Y. et al. Lipids 29(3): 189-193, 1994.
16. Yeh, Y. Shinyaku To Rinsho (New Drug Clin.) 45(3): 441-450, 1996.
17. Yeh, Y. et al. In: Food Factors for Cancer Prevention. Ohigashi, et al. eds. Springer-ag, Tokyo, p 226-230, 1997.
18. Kikuchi, T. et al. Shinyaku To Rinsho (Jpn. J. New Remedies Clin.) 43(1): 146-158, 1994.
19. Okuhara, T. Jpn. Pharmacol. Therapeut. 22(8): 3695-3701, 1994.
20. Steiner, M. et al. J. Cardiovascular Pharmacol. 31: 904-908, 1998.
21. Yokoyama, K. et al. Oyo Yakuri (Appl. Pharmacol.) 36: 301-308, 1988.
22. Abdullah, T. et al. J. Oncology 21: 52-53, 1989.
23. Hirao, Y. et al. Phytother. Res. 1: 161-164, 1987.
24. Kandil, O. et al. Arch. AIDS Res 1:230-231, 1988.
25. Kyo, E. et al. Phytomed. 5(4): 259-267, 1998.
26. Lau, B.H.S. et al. 1991. Molecular Biother. 3: 103-107.
27. Lau, B.H.S. Int. Clin. Nutr. Rev. 9: 27-31, 1989.
28. Morioka, N. et al. Cancer Immunol. Immunother. 37: 316-322, 1993.
29. Nagai, K. 1973. Kansenshogaku-Zasshi 47(9): 321-325.
30. Reeve, V. et al. Photochem. Photobiol. 58(6): 813-817, 1993.
31. Riggs, D. et al. Cancer 79: 1987-1994, 1997.
32. Tadi, P.P. et al. Int. Clin. Nutr. Rev. 10: 423-429, 1990.
33. Hasegawa, Y. et al. Shinyaku To Rinsho (Jpn. J. New Remedies Clin.) 32: 365-376, 1983.
34. Kawashima, H. et al. Shinryou To Shinyaku (Treat. New Med.) 22(12): 3012-3024, 1985.

35. Kawashima, Y. et al. Kiso To Rinsho (Preclin. Clin. Reports) 20(16): 8229-8245, 1984.
36. Miyoshi, A. et al. Shinryou To Shinyaku (Treat. New Med.) 21: 1806-1820, 1984.
37. Okada, K. and Miyagaki, H. Kiso To Rinsho (Preclin. Clin. Reports) 17: 2173-2183, 1983.
38. Takasugi, N. et al. Oyo Yakuri (Appl. Pharmacol.) 31: 967-976, 1986.
39. Takasugi, N. et al. Oyo Yakuri (Appl. Pharmacol.) 28: 991-1002, 1984.
40. Yokoyama, K. et al. Oyo Yakuri (Appl. Pharmacol.) 31: 977-984, 1986.
41. Horie, T. et al. Planta Med. 55: 506-508, 1989.
42. Horie, T. et al. Planta Med. 58: 468-469, 1992.
43. Nakagawa, S. et al. Hiroshima J. Med. Sci. 34: 303-309, 1985.
44. Nakagawa, S. et al. Phytother. Res. 1: 1-4, 1988.
45. Sumioka, I. et al. Jpn. J. Pharmacol. 78: 199-207, 1998.
46. Tadi, P. et al. Nutr. Cancer 15: 87-95, 1991
47. Wang, B. et al. Toxicology 126: 213-222, 1998.
48. Wang, B. et al. Toxicology 132: 215-225, 1999.
49. Welch, C. et al. Cancer Lett. 63: 211-219, 1992.
50. Gwebu, E. et al. Soc. Neurosci. 24(2):#571.11, 1998.
51. Moriguchi, T. et al. Neurochem. Res. 22(12): 1449-1452, 1997.
52. Moriguchi, T. et al. Phytother. Res. 10: 468-472, 1996.
53. Nishiyama, N. et al. Int. Acad. Biomed. Drug Res. Basel. Karger. 11: 253-258, 1996.
54. Saito, H. et al. Ch. 16. In: Nutraceuticals: Designer Foods III Garlic, Soy and Licorice. Paul P. Lanchance (ed.). Food & Nutrition Press, Trumbell, CT, pp. 153-161, 1997.
55. Matsuura, H. et al. Chem. Pharm. Bull. 36: 3659-3663, 1988.
56. Recommended Dietary Allowances (RDA). 10th ed. National Academy Press, Washington, D.C., 1989.
57. Kokai Tokyo Koho, Japanese Patent H1-252276.
58. Matsuura, H. Phytochemistry of Garlic Horticultural and Processing Procedures. Designer Foods III Symposium: Research Update on Phytochemicals in Garlic, Soy and Licorice in Washington, D.C., May 1994.
59. Forsum, E. J. Dairy Sci. 57(6): 665-670, 1974.
60. Petschow, B. & Talbott, R. Clin. Microbiol. 28(2): 287-292, 1990.

