

- Contains Aged Garlic Extract™ [300 mg], Vitamin B6 [12.5mg], Vitamin B12 [100mcg], Folic acid [200mcg], Trimethylglycine [125mg] (per capsule)
- Studies suggest that Aged Garlic Extract and its constituents may provide nutritional support for healthy cardiovascular function, including homocysteine, cholesterol, blood pressure, and stimulating circulation, all while inhibiting oxidative damage in the body.[1-20]
- Kyolic is already patented for its ability to reduce homocysteine levels, now thought to be a major risk factor for heart disease.[27,28]
- In a study of 734 subjects, homocysteine levels were found to be highest in subjects with the lowest intake of folic acid, vitamin B12 and vitamin B6[22].
- Trimethylglycine has been reported to reduce blood homocysteine levels, since it plays a role as a methyl donor, helping to detoxify homocysteine in the blood.[23-26]
- Vitamin B6 and B12, Folic Acid and Trimethylglycine (as TMG or Betaine) are added since their synergy is found more effective than any one ingredient alone in facilitating the proper function of homocysteine recycling (detoxification)

- **Supplement Facts**
- **Scientific References**
- **Next Kyolic Product**
- **Previous Kyolic Product**
- **All Kyolic Products**
- **Consumer Price List**

Kyolic Formula 108

SUGGESTED USE:
Take two capsules **with a meal** twice daily.

Supplement Facts Serving Size 2 capsules		**Product Code:** **108-41: 100 capsules**
	Amount per serving	%Daily Value
Aged Garlic Extract™ Powder (bulb)	500mg	**
Vitamin B6	25mg	1250%
Folic Acid	400mcg	100%
Vitamin B12	200mcg	3333%

Trimethylglycine (TMG)(as Betaine)	250mg	**

**Daily Value not established

OTHER INGREDIENTS:
Gelatin, Cellulose, Magnesium Stearate (vegetable source) and Silica.

Free of: gluten, preservatives, sugar, sodium, milk, yeast, artificial colors and flavors.



REFERENCES FOR KYOLIC Formula 108

1. Yeh, Y. Y. et al. 1994. Lipids 29(3): 189-193.
2. Yeh, Y. Y. et al. 1990. First World Congress on the Health Significance of Garlic and Garlic Constituents. Nutrition International Co., Irvine, CA., p. 37.
3. Kawashima, Y. et al. 1989. Shinryo To Shinyaku (Treat. New Med.) 26: 377-388.
4. Lau, B. et al. 1987. Nutr. Res. 7: 139-149.
5. Steiner, M. et al. 1994. J. Am. Coll. Nutr. 13(5): 524.
6. Yeh, Y. et al. 1995. J. Am. Coll. Nutr. 14(5):545.
7. Abuirmeileh, N. et al. 1991. FASEB J. 5(6): 8048.
8. Qureshi, N. et al. 1990. First World Congress on the Health Significance of Garlic and Garlic Constituents. Nutrition International Co., Irvine, CA, p. 17.
9. Qureshi, A. et al. 1990. First World Congress on the Health Significance of Garlic and Garlic Constituents. Nutrition International Co., Irvine, CA, p. 16.
10. Yeh, Y.Y. 1996. Shinyaku To Rinsho (New Drug Clin.) 45(3): 441-450.
11. Efendy, J.L.et al.1996. In: Proceedings of the 1996 Conference of the Anatomical Society of Australia and New Zealand. The Department of Anatomical Sciences, the University of Queensland. Brisbane, Australia, p. 19.
12. Efendy, J.L. et al. 1996. J. Vasc. Res. 33(S1): 23.
13. Steiner, M. et al.1996. 2nd International Congress on Phytomedicine. September 11-14, 1996. Munich, Germany.
14. Kawashima, Y. et al. 1989. Shinryou to shinyaku (Treat. New Med.) 26: 377-388.
15. Okuhara, T. 1994. Jpn. Pharmacol. Therapeut. 22(8): 3695-3701.
16. Kikuchi, T. et al. 1994. Shinyaku To Rinsho (Jpn. J. New Remedies Clin.) 43(1): 146-158.
17. Kawashima, Y. et al. Shinryou to shinyaku (Treat. New Med.) 26: 377-388, 1989.
18. Okuhara, T. 1994. Jpn. Pharmacol. Therapeut. 22(8): 3695-3701.
19. Kikuchi, T. et al. Shinyaku To Rinsho (Jpn. J. New Remedies Clin.) 43(1): 146-158, 1994.
20. Steiner, M. et al. Am. J. Clin. Nutr. 64: 866-870, 1996.
21. McCully, K. JAMA, 279(5): 392-395. 1998
22. Darlery K, et al. Am. J. Cardiol. 75(16): 1107-1111. 1995
23. Selhub J. Annu Rev Nutr 1999;19:217-46 [review].
24. Wendel U, Bremer HJ. Eur J Pediatr 1984;142:147-50.
25. Wilcken DE, Wilcken B, Dudman NP, Tyrrell PA. N Engl J Med 1983;309:448-53.
26. Gahl WA, Bernardini I, Chen S, et al. J Inherit Metab Dis 1988;11:291-8.
27. Yeh, Y., et al. 1999 FASEB 13(4): a232; #209.12
28. Amagase, H. 2000. Method and pharmaceutical composition for reducing serum homocysteine concentrations. Patent # 6,129,918.

